arXiv:0806.2431v4 [math.AG] 27 Jul 2009

Polynomial Hamiltonian system in two variables with $W(A_1^{(1)})$-symmetry and the second member of the second Painlevé hierarchy

By
Yusuke Sasano

July 27, 2009

Abstract

We find a one-parameter family of polynomial Hamiltonian system in two variables with $W(A_1^{(1)})$-symmetry. We also show that this system can be obtained by the compatibility conditions for the linear differential equations in three variables. We give a relation between it and the second member of the second Painlevé hierarchy.

Key Words and Phrases. Bäcklund transformation, Birational transformation, Holomorphy condition, Painlevé equations, Second Painlevé hierarchy.

2000 *Mathematics Subject Classification Numbers.* 34M55; 34M45; 58F05; 32S65.

1 Introduction

In this paper, we find a 1-parameter family of total differential system in two variables t, s

$$\begin{cases} dx =(-2xp-\alpha_0)dt+(-2xz-2\alpha_0 w)ds,\\ dy =(2yp+\alpha_1)dt+(2yz-2\alpha_1 q)ds,\\ dz =\left(\dfrac{x+y}{2}\right)dt+(-xq+yw-2p)ds,\\ dw =(z-2wp)dt+(-2xp-2\alpha_0-\alpha_1)ds,\\ dq =(z+2qp)dt+(2yp+\alpha_0+2\alpha_1)ds,\\ dp =\left(\dfrac{w+q}{2}\right)dt+\left(\dfrac{x+y}{2}\right)ds, \end{cases} \tag{1}$$

where the constant parameters α_i satisfy the relation $\alpha_0+\alpha_1=1$.
In next theorem, we show that this system satisfies the compatibility conditions, and we can characterize this system by the following holomorphy conditions r_0, r_1.

Theorem 1.1. *Let us consider a system of first order ordinary differential equations in the polynomial class*:

$$\frac{dx}{dT}=f_1(x,y,z,w,q,p),\dots,\frac{dp}{dT}=f_6(x,y,z,w,q,p) \quad f_i\in\mathbb{C}[x,y,z,w,q,p] \quad (i=1,\dots,6).$$

We assume that
$(A1)$ $deg(f_i)=2$ with respect to x,y,z,w,q,p.
$(A2)$ The right-hand side of this system becomes again a polynomial in each coordinate system r_i $(i=0,1)$:

$$r_0: x_0=-(xp+\alpha_0)p, \quad y_0=y-4zp+4wp^2, \quad z_0=z-2wp, \quad w_0=w, \quad q_0=q-2p^2, \quad p_0=\frac{1}{p},$$

$$r_1: x_1=x+4zp+4qp^2, \quad y_1=-(yp+\alpha_1)p, \quad z_1=z+2qp, \quad w_1=w+2p^2, \quad q_1=q, \quad p_1=\frac{1}{p}.$$

Then we can obtain two systems:

$$\begin{cases} \dfrac{dx}{dt} =& -2xp - \alpha_0, \\ \dfrac{dy}{dt} =& 2yp + \alpha_1, \\ \dfrac{dz}{dt} =& \dfrac{x+y}{2}, \\ \dfrac{dw}{dt} =& z - 2wp, \\ \dfrac{dq}{dt} =& z + 2qp, \\ \dfrac{dp}{dt} =& \dfrac{w+q}{2} \end{cases} \tag{2}$$

and

$$\begin{cases} \dfrac{dx}{ds} =& -2xz - 2\alpha_0 w, \\ \dfrac{dy}{ds} =& 2yz - 2\alpha_1 q, \\ \dfrac{dz}{ds} =& 2yz - 2\alpha_1 q, \\ \dfrac{dw}{ds} =& -2xp - 2\alpha_0 - \alpha_1, \\ \dfrac{dq}{ds} =& 2yp + \alpha_0 + 2\alpha_1, \\ \dfrac{dp}{ds} =& \dfrac{x+y}{2}. \end{cases} \tag{3}$$

These two systems satisfy the compatibility conditions:

$$\frac{\partial}{\partial s}\frac{\partial x}{\partial t} = \frac{\partial}{\partial t}\frac{\partial x}{\partial s}, \quad \frac{\partial}{\partial s}\frac{\partial y}{\partial t} = \frac{\partial}{\partial t}\frac{\partial y}{\partial s}, \dots, \frac{\partial}{\partial s}\frac{\partial p}{\partial t} = \frac{\partial}{\partial t}\frac{\partial p}{\partial s}. \tag{4}$$

We remark that these transition functions in r_0, r_1 satisfy the condition:

$$dx_i \wedge dy_i \wedge dz_i \wedge dw_i \wedge dq_i \wedge dp_i = dx \wedge dy \wedge dz \wedge dw \wedge dq \wedge dp \quad (i = 0, 1). \tag{5}$$

Theorem 1.2. *The system* (1) *admits the extended affine Weyl group symmetry of type* $A_1^{(1)}$ *as the group of its Bäcklund transformations, whose generators* s_0, s_1, π *defined as follows*: *with the notation* $(*) := (x, y, z, w, q, p; \alpha_0, \alpha_1)$:

$$\begin{aligned} s_0 : (*) &\rightarrow \left(x, y + \frac{4\alpha_0 z}{x} + \frac{4\alpha_0^2 w}{x^2}, z + \frac{2\alpha_0 w}{x}, w, q + \frac{4\alpha_0 p}{x} + \frac{2\alpha_0^2}{x^2}, p + \frac{\alpha_0}{x}; -\alpha_0, \alpha_1 + 2\alpha_0\right), \\ s_1 : (*) &\rightarrow \left(x - \frac{4\alpha_1 z}{y} + \frac{4\alpha_1^2 q}{y^2}, y, z - \frac{2\alpha_1 q}{y}, w - \frac{4\alpha_1 p}{y} - \frac{2\alpha_1^2}{y^2}, q, p + \frac{\alpha_1}{y}; \alpha_0 + 2\alpha_1, -\alpha_1\right), \\ \pi : (*) &\rightarrow (-y, -x, -z, -q, -w, -p; \alpha_1, \alpha_0). \end{aligned}$$

2 Lax pair for the system

In this section, we show that the system (1) can be obtained by the compatibility conditions for the linear differential equations of second order in three variables T, t, s

$$T\partial_T \vec{u} = A(T, t)\vec{u}, \quad \partial_t \vec{u} = B_1(T, t)\vec{u}, \quad \vec{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}, \tag{6}$$

and

$$T\partial_T \vec{u} = A(T, s)\vec{u}, \quad \partial_s \vec{u} = B_3(T, s)\vec{u}. \tag{7}$$

The matrices $A(T,t), B_1(T,t)$ and $B_3(T,s)$ are explicitly given by

$$\begin{aligned}
-A(T,t) &= \begin{pmatrix} \varepsilon_1 & y \\ 0 & \varepsilon_2 \end{pmatrix} + \begin{pmatrix} 2z & 4q \\ -x & -2z \end{pmatrix} T + \begin{pmatrix} 8p & -8 \\ -4w & -8p \end{pmatrix} T^2 + \begin{pmatrix} 0 & 0 \\ -8 & 0 \end{pmatrix} T^3, \\
B_1(T,t) &= \begin{pmatrix} p & -1 \\ 0 & -p \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix} T, \\
B_3(T,s) &= \begin{pmatrix} \frac{z}{4} & \frac{q}{2} \\ 0 & -\frac{z}{4} \end{pmatrix} + \begin{pmatrix} p & -1 \\ -\frac{w}{2} & -p \end{pmatrix} T + \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix} T^2.
\end{aligned} \tag{8}$$

Here, the matrices $A(T,t), B_1(T,t)$ and $B_3(T,s)$ depend on t, s, and ε_i are constant parameters. At first, by the compatibility conditions

$$\partial_t(A(T,t)) - T\partial_T(B_1(T,t)) + [A(T,t), B_1(T,t)] = 0 \tag{9}$$

for the linear differential equations in two variables T, t

$$T\partial_T \vec{u} = A(T,t)\vec{u}, \quad \partial_t \vec{u} = B_1(T,t)\vec{u}, \tag{10}$$

we can obtain the non-linear ordinary differential system in the variable t

$$\begin{cases}
\dfrac{dx}{dt} = -2xp - 1 + \varepsilon_1 - \varepsilon_2, \\
\dfrac{dy}{dt} = 2yp + \varepsilon_1 - \varepsilon_2, \\
\dfrac{dz}{dt} = \dfrac{x+y}{2}, \\
\dfrac{dw}{dt} = -2wp + z, \\
\dfrac{dq}{dt} = 2qp + z, \\
\dfrac{dp}{dt} = \dfrac{w+q}{2},
\end{cases} \tag{11}$$

where ε_i satisfy the relations

$$\begin{cases}
\alpha_0 = -\varepsilon_1 + \varepsilon_2 + 1, \\
\alpha_1 = \varepsilon_1 - \varepsilon_2.
\end{cases} \tag{12}$$

Next, by the compatibility conditions

$$\partial_s(A(T,s)) - T\partial_T(B_3(T,s)) + [A(T,s), B_3(T,s)] = 0 \tag{13}$$

for the linear differential equations in two variables T, s

$$T\partial_T \vec{u} = A(T,s)\vec{u}, \quad \partial_s \vec{u} = B_3(T,s)\vec{u}, \tag{14}$$

we can obtain the non-linear ordinary differential system in the variable s

$$\begin{cases}
\dfrac{dx}{ds} = -2xz - 2(1 - \varepsilon_1 + \varepsilon_2)w, \\
\dfrac{dy}{ds} = 2yz - 2(\varepsilon_1 - \varepsilon_2)q, \\
\dfrac{dz}{ds} = -xq + yw - 2p, \\
\dfrac{dw}{ds} = -2xp - 2(1 - \varepsilon_1 + \varepsilon_2) - (\varepsilon_1 - \varepsilon_2), \\
\dfrac{dq}{ds} = 2yp + (1 - \varepsilon_1 + \varepsilon_2) + 2(\varepsilon_1 - \varepsilon_2), \\
\dfrac{dp}{ds} = \dfrac{x+y}{2}.
\end{cases} \tag{15}$$

3 Polynomial Hamiltonian system

In this section, we show that the system (1) is equivalent to the polynomial Hamiltonian system in two variables t, s. At first, we show that the system (1) has two first integrals.

Proposition 3.1. *This system* (1) *has its first integrals:*

$$\begin{cases} w - q + 2p^2 + 3s = C_1, \\ 8sp^2 + 4zp - 2wq + x - y + 4sw - 4sq + t + 6s^2 = C_2, \quad (C_1, C_2 \in \mathbb{C}). \end{cases} \tag{16}$$

Theorem 3.1. *The transformations*

$$\begin{cases} q_1 = p, \\ p_1 = x, \\ q_2 = z - 2wp, \\ p_2 = w \end{cases} \tag{17}$$

take the system (1) *to the polynomial Hamiltonian system in two variables* t, s

$$\begin{cases} dq_1 = \dfrac{\partial H_1}{\partial p_1} dt + \dfrac{\partial H_2}{\partial p_1} ds, \\ dp_1 = -\dfrac{\partial H_1}{\partial q_1} dt - \dfrac{\partial H_2}{\partial q_1} ds, \\ dq_2 = \dfrac{\partial H_1}{\partial p_2} dt + \dfrac{\partial H_2}{\partial p_2} ds, \\ dp_2 = -\dfrac{\partial H_1}{\partial q_2} dt - \dfrac{\partial H_2}{\partial q_2} ds \end{cases} \tag{18}$$

with the polynomial Hamiltonians (cf. [2])

$$\begin{aligned} H_1 =& q_1^2 p_1 + \frac{3}{2} s p_1 - \frac{C_1}{2} p_1 + \alpha_0 q_1 - p_2^3 + (C_1 - 3s) p_2^2 + \left(-3s^2 + 2C_1 s + \frac{t}{2} - \frac{C_2}{2} \right) p_2 - \frac{q_2^2}{2} + p_1 p_2, \\ H_2 =& \frac{p_1^2}{2} - \frac{1}{2}(C_2 - 4C_1 s + 6s^2 - t) p_1 + (2\alpha_0 + \alpha_1) q_2 \\ & - p_1 p_2^2 + 2q_1^2 p_1 p_2 + 2q_1 p_1 q_2 + 2\alpha_0 q_1 p_2 + (C_1 - 3s) p_1 p_2. \end{aligned} \tag{19}$$

Recently, H. Kawamuko showed that this system can be obtained by the monodromy preserving deformations of half-integer type equation $L(7/2, 1)$ (see [4]).
The relations between x, y, z, w, q, p and q_1, p_1, q_2, p_2 are given by

$$\begin{cases} x = p_1, \\ y = 4q_1^2 p_2 - 2p_2^2 + 4q_1 q_2 + 2C_1 p_2 - 6s p_2 + p_1 - 6s^2 + t + 4C_1 s - C_2, \\ z = q_2 + 2q_1 p_2, \\ w = p_2, \\ q = 2q_1^2 + p_2 + 3s - C_1, \\ p = q_1. \end{cases} \tag{20}$$

Setting $C_1 = 0, C_2 = 0, s = 0$ and $\alpha_0 := \frac{1}{2} - \alpha_2$ in the Hamiltonian H_1, we can obtain the Hamiltonian in the variable t given by [5]. This Hamiltonian system is equivalent to the second member of the second Painlevé hierarchy (see [5])

$$P_{II}^{(2)} : \frac{d^4 u}{dt^4} = 10u \left(\frac{du}{dt} \right)^2 + 10u^2 \frac{d^2 u}{dt^2} - 6u^5 + tu + \alpha_2 \quad (\alpha_2 \in \mathbb{C}). \tag{21}$$

4 Symmetry and holomorphy of the system (18)

Theorem 4.1. *The system* (18) *admits the extended affine Weyl group symmetry of type* $A_1^{(1)}$ *as the group of its Bäcklund transformations, whose generators* s_0, s_1, π *defined as follows*: *with the notation* $(*) := (q_1, p_1, q_2, p_2, t, s; \alpha_0, \alpha_0)$:

$$
\begin{aligned}
s_0 : (*) \rightarrow &\left(q_1 + \frac{\alpha_0}{f_0}, p_1, q_2, p_2, t, s; -\alpha_0, \alpha_1 + 2\alpha_0\right), \\
s_1 : (*) \rightarrow &(q_1 + \frac{\alpha_1}{f_1}, p_1 - \frac{4\alpha_1(q_2 + 2q_1p_2)}{f_1} + \frac{4\alpha_1^2(p_2 + 2q_1^2 - (C_1 - 3s))}{f_1^2}, \\
&q_2 - \frac{2\alpha_1(2p_2 - 2q_1^2 - (C_1 - 3s))}{f_1} + \frac{12\alpha_1^2 q_1}{f_1^2} + \frac{4\alpha_1^3}{f_1^3}, p_2 - \frac{4\alpha_1 q_1}{f_1} - \frac{2\alpha_1^2}{f_1^2}, t, s; \alpha_0 + 2\alpha_1 - \alpha_1), \\
\pi : (*) \rightarrow &(-q_1, -f_1, -(q_2 + 4q_1(q_1^2 + p_2) - 2(C_1 - 3s)q_1), -(p_2 + 2q_1^2 - C_1 + 3s), t, s; \alpha_1, \alpha_0),
\end{aligned}
$$

where $f_0 := p_1$ and $f_1 := p_1 + 4q_1^2p_2 - 2p_2^2 + 4q_1q_2 + 2(C_1 - 3s)p_2 - C_2 + 4C_1s - 6s^2 + t$.

We note that the Bäcklund transformations of this system satisfy the universal description for $A_1^{(1)}$ root system:

$$
s_i(g) = g + \frac{\alpha_i}{f_i}\{f_i, g\} + \frac{1}{2!}\left(\frac{\alpha_i}{f_i}\right)^2 \{f_i, \{f_i, g\}\} + \cdots \quad (g \in \mathbb{C}(t, s)[q_1, p_1, q_2, p_2]), \tag{22}
$$

where $\{p_i, q_j\} = \delta_{ij}$ and $\{p_i, p_j\} = \{q_i, q_j\} = 0$.

Theorem 4.2. *Let us consider a polynomial Hamiltonian system with Hamiltonian* $H \in \mathbb{C}(t, s)[q_1, p_1, q_2, p_2]$. *We assume that*
(A1) $deg(H) = 5$ *with respect to* q_1, p_1, q_2, p_2.
(A2) This system becomes again a polynomial Hamiltonian system in each coordinate $\tilde{r}_i$ $(i = 0, 1)$:

$$
\begin{aligned}
\tilde{r}_0 : (x_0, y_0, z_0, w_0) &= \left(\frac{1}{q_1}, -(q_1 f_0 + \alpha_0)q_1, q_2, p_2\right), \\
\tilde{r}_1 : (x_1, y_1, z_1, w_1) &= \left(\frac{1}{q_1}, -(q_1 f_1 + \alpha_1)q_1, q_2 + 4q_1(q_1^2 + p_2) - 2(C_1 - 3s)q_1, p_2 + 2q_1^2\right).
\end{aligned}
$$

Then such a system coincides with the Hamiltonian system (18) *with the polynomial Hamiltonians* H_1, H_2.

We note that the conditions $(A2)$ should be read that

$$
\begin{aligned}
&r_0(H), \quad r_0(H - q_1), \\
&r_1(H), \quad r_1(H + 8q_1^3 + 6q_1p_2 - 4(C_1 - 3s)q_1)
\end{aligned}
$$

are polynomials with respect to q_1, p_1, q_2, p_2.

5 Particular solutions

In this section, we study a solution of the system (18) which is written by the use of known functions. By the transformation π, the fixed solution is derived from

$$
\begin{aligned}
&\alpha_0 = \alpha_1, \quad q_1 = -q_1, \quad p_1 = -f_1, \\
&q_2 = -(q_2 + 4q_1(q_1^2 + p_2) - 2(C_1 - 3s)q_1), \quad p_2 = -(p_2 + 2q_1^2 - C_1 + 3s).
\end{aligned} \tag{23}
$$

Then we obtain

$$
(q_1, p_1, q_2, p_2; \alpha_0, \alpha_1) = \left(0, \frac{-2t + 3s^2 - 2C_1s - C_1^2 + 2C_2}{4}, 0, \frac{C_1 - 3s}{2}; \frac{1}{2}, \frac{1}{2}\right) \tag{24}
$$

as a seed solution.

References

[1] P. Clarkson, N. Joshi and A. Pickering, *Bäcklund transformations for the second Painlevé hierarchy: a modified truncation approach*, Inverse Problem. **15** (1999), 175–187.

[2] M. Mazzocco and M. Y. Mo, *The Hamiltonian structure of the second Painlevé hierarchy*, http://arXiv.org/abs/nlin.SI/0610066.

[3] N. Joshi, *The second Painlevé hierarchy and the stationary KdV hierarchy*, Publ. RIMS. Kyoto Univ. **40** (2004), 1039-1061.

[4] H. Kawamuko, *On the Garnier system of half-integer type in two variables*, to appear in Funkcial. Ekvac..

[5] Y. Sasano, *Symmetry and holomorphy of the second member of the second Painlevé hierarchy*, preprint.